COMPONENTS:

- (1) Acetamide, N-[4-[[(4-methoxy-1,2,5thiadiazo1-3-y1)amino]sulfony1]pheny1]-(N⁴-acetylsulfametrole); C₁₁H₁₂N₄O₄S₂; [79962-97-3] Phosphoric acid, disodium salt;
- (2) Na₂HPO₄; [7558-94-4]
- (3) Phosphoric acid, monopotassium salt; KH₂PO₄; [7778-77-0]
- (4) Water; H₂0; [7732-18-5]

VARIABLES:

рΗ

ORIGINAL MEASUREMENTS:

Hekster, Y. A.; Vree, T. B.; Damsma, J. E.; Friesen, W. T. J. Antimicrob. Chemother. 1981, 8. 133-44.

PREPARED BY:

R. Piekos

EXPERIMENTAL VALUES:

рН	Solubility at 25°C	
	mg/l	10^3 mol dm ⁻³ a
5.5	1100	3.350
7.5	6000	18.273

^aCalculated by compiler

AUXILIARY INFORMATION

METHOD/APPARATUS/PROCEDURE:

Satd solns of N4-acetylsulfametrole were prepd in phosphate buffers of pH 5.5 and 7.5 at room temp (25°C). The concn of the solute was measured by means of a Spectra Physics 3500B high-performance liquid chromatograph equipped with a column oven (Model 748) and a Pye-Unicam LC-UV spectrophotometric detector. The detector was connected to a 1-mV recorder. A stainless steel column (10 cm x 4.6 mm i.d.) was packed with Lichrosorb RPS, 5 μm, obtained from Chrompack. An injection loop of 100 μ l was used. The oven temp was 40°C. Detection of the solute was performed at 260 nm.

SOURCE AND PURITY OF MATERIALS:

 N^4 -acetylsulfametrole was obtained from Chemie, Linz. The compd was 100% pure according to the HPLC chromatogram. The source and purity of the remaining materials were not specified.

ESTIMATED ERROR:

The detection limit of the solute by HPLC was 0.5 mg/1 (authors). The error in temperature and pH were not specified.

REFERENCES: